

**(19) World Intellectual Property
Organization
International Bureau**



(43) International Publication Date
23 June 2005 (23.06.2005)

PCT

(10) International Publication Number
WO 2005/056530 A1

(51) International Patent Classification⁷: C07D 213/80,
A61K 31/455, A61P 03/04, 03/10

(21) International Application Number:
PCT/GB2004/005068

SK10 4TG (GB). MCKERRECHER, Darren [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). **PIKE, Kurt, Gordon [GB/GB];** AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(22) International Filing Date: 2 December 2004 (02.12.2004)

(74) Agent: ASTRAZENECA; Global Intellectual Property, SE-151 85 Sodertalje (SE).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0328178.9 5 December 2003 (05.12.2003) GB

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) **Applicant** (for AE, AG, AL, AM, AT, AU, AZ, BA, BB, BE, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CY, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MN, MW, MX, MC, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, SZ, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW only): **ASTRAZENECA AB** [SE/SE]: SE-151 85 Sodertalje (SE).

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).

(72) Inventors: and


Published:

(75) **Inventors/Applicants (for US only): CAULKETT, Peter, William, Rodney [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). JOHNSTONE, Craig [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire**

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BENZOYL AMINO PYRIDYL CARBOXYLIC ACID DERIVATIVES USEFUL AS GLUCOKINASE (GLK) ACTIVATORS



(57) **Abstract:** A compound of Formula (I): Formula (I) wherein: R¹ is selected from hydrogen and C₁₋₄alkyl; R² is selected from: R⁴-C(R^{5a}R^{5b})-, R⁴=C(R⁶)- and R^{7a}C(R^{7b})=C(R⁶)-; R³ X- is selected from methyl, methoxymethyl and ; R⁴ is selected from (optionally substituted) C₁₋₄alkyl, phenyl, C₃₋₆cycloalkyl and heteroaryl; R^{5a} and R^{5b} are independently selected from hydrogen, fluoro and C₁₋₄alkyl; R⁶ is selected from hydrogen and C₁₋₄alkyl; R^{7a} and R^{7b} are optionally substituted C₁₋₄alkyl; or a salt, pro-drug or solvate thereof, are described. Their use as GLK activators, pharmaceutical compositions containing them, and processes for their preparation are also described.

WO 2005/056530 A1

BENZOYL AMINO PYRIDYL CARBOXYLIC ACID DERIVATIVES USEFUL AS GLUCOKINASE (GLK) ACTIVATORS

The present invention relates to a group of benzoyl amino pyridyl carboxylic acids which are useful in the treatment or prevention of a disease or medical condition mediated through glucokinase (GLK), leading to a decreased glucose threshold for insulin secretion. In addition the compounds are predicted to lower blood glucose by increasing hepatic glucose uptake. Such compounds may have utility in the treatment of Type 2 diabetes and obesity. The invention also relates to pharmaceutical compositions comprising said compounds and to methods of treatment of diseases mediated by GLK using said compounds.

10 In the pancreatic β -cell and liver parenchymal cells the main plasma membrane glucose transporter is GLUT2. Under physiological glucose concentrations the rate at which GLUT2 transports glucose across the membrane is not rate limiting to the overall rate of glucose uptake in these cells. The rate of glucose uptake is limited by the rate of phosphorylation of glucose to glucose-6-phosphate (G-6-P) which is catalysed by glucokinase (GLK) [1]. GLK has a high (6-10mM) K_m for glucose and is not inhibited by physiological concentrations of G-6-P [1]. GLK expression is limited to a few tissues and cell types, most notably pancreatic β -cells and liver cells (hepatocytes) [1]. In these cells GLK activity is rate limiting for glucose utilisation and therefore regulates the extent of glucose induced insulin secretion and hepatic glycogen synthesis. These processes are critical in the maintenance of whole body glucose homeostasis and both are dysfunctional in diabetes [2].

In one sub-type of diabetes, Type 2 maturity-onset diabetes of the young (MODY-2), the diabetes is caused by GLK loss of function mutations [3, 4]. Hyperglycaemia in MODY-2 patients results from defective glucose utilisation in both the pancreas and liver [5]. Defective glucose utilisation in the pancreas of MODY-2 patients results in a raised threshold for glucose stimulated insulin secretion. Conversely, rare activating mutations of GLK reduce this threshold resulting in familial hyperinsulinism [6, 6a, 7]. In addition to the reduced GLK activity observed in MODY-2 diabetics, hepatic glucokinase activity is also decreased in type 2 diabetics [8]. Importantly, global or liver selective overexpression of GLK prevents or reverses the development of the diabetic phenotype in both dietary and genetic models of the disease [9-12]. Moreover, acute treatment of type 2 diabetics with fructose improves glucose tolerance through stimulation of hepatic glucose utilisation [13]. This effect is believed to be

mediated through a fructose induced increase in cytosolic GLK activity in the hepatocyte by the mechanism described below [13].

Hepatic GLK activity is inhibited through association with GLK regulatory protein (GLKRP). The GLK/GLKRP complex is stabilised by fructose-6-phosphate (F6P) binding to
5 the GLKRP and destabilised by displacement of this sugar phosphate by fructose-1-phosphate (F1P). F1P is generated by fructokinase mediated phosphorylation of dietary fructose. Consequently, GLK/GLKRP complex integrity and hepatic GLK activity is regulated in a nutritionally dependent manner as F6P is elevated in the post-absorptive state whereas F1P predominates in the post-prandial state. In contrast to the hepatocyte, the pancreatic β -cell
10 expresses GLK in the absence of GLKRP. Therefore, β -cell GLK activity is regulated exclusively by the availability of its substrate, glucose. Small molecules may activate GLK either directly or through destabilising the GLK/GLKRP complex. The former class of compounds are predicted to stimulate glucose utilisation in both the liver and the pancreas whereas the latter are predicted to act exclusively in the liver. However, compounds with
15 either profile are predicted to be of therapeutic benefit in treating Type 2 diabetes as this disease is characterised by defective glucose utilisation in both tissues.

GLK and GLKRP and the K_{ATP} channel are expressed in neurones of the hypothalamus, a region of the brain that is important in the regulation of energy balance and the control of food intake [14-18]. These neurones have been shown to express orectic and
20 anorectic neuropeptides [15, 19, 20] and have been assumed to be the glucose-sensing neurones within the hypothalamus that are either inhibited or excited by changes in ambient glucose concentrations [17, 19, 21, 22]. The ability of these neurones to sense changes in glucose levels is defective in a variety of genetic and experimentally induced models of obesity [23-28]. Intracerebroventricular (icv) infusion of glucose analogues, that are
25 competitive inhibitors of glucokinase, stimulate food intake in lean rats [29, 30]. In contrast, icv infusion of glucose suppresses feeding [31]. Thus, small molecule activators of GLK may decrease food intake and weight gain through central effects on GLK. Therefore, GLK activators may be of therapeutic use in treating eating disorders, including obesity, in addition to diabetes. The hypothalamic effects will be additive or synergistic to the effects of the same
30 compounds acting in the liver and/or pancreas in normalising glucose homeostasis, for the treatment of Type 2 diabetes. Thus the GLK/GLKRP system can be described as a potential "Diabetes" target (of benefit in both Diabetes and Obesity).

In WO0058293 and WO01/44216 (Roche), a series of benzylcarbamoyl compounds are described as glucokinase activators. The mechanism by which such compounds activate GLK is assessed by measuring the direct effect of such compounds in an assay in which GLK activity is linked to NADH production, which in turn is measured optically - see details of the
5 *in vitro* assay described in Example A. Many compounds of the present invention may show favourable selectivity compared to known GLK activators.

WO9622282, WO9622293, WO9622294, WO9622295, WO9749707 and WO9749708 disclose a number of intermediates used in the preparation of compounds useful as vasopressin agents which are structurally similar to those disclosed in the present invention.
10 Structurally similar compounds are also disclosed in WO9641795 and JP8143565 (vasopressin antagonism), in JP8301760 (skin damage prevention) and in EP619116 (osetopathy).

WO01/12621 describes the preparation of as isoxazolympyrimidines and related compounds as inhibitors of c-JUN N-terminal kinases, and pharmaceutical compositions
15 containing such compounds.


Cushman *et al* [Bioorg Med Chem Lett (1991) 1(4), 211-14] describe the synthesis of pyridine-containing stilbenes and amides and their evaluation as protein-tyrosine kinase inhibitors. Rogers *et al* [J Med Chem (1981) 24(11) 1284-7] describe mesoionic purinone analogs as inhibitors of cyclic-AMP phosphodiesterase.

20 WO00/26202 describes the preparation of 2-amino-thiazole derivatives as antitumour agents. GB 2331748 describes the preparation of insecticidal thiazole derivatives. WO96/36619 describes the preparation of aminothiazole derivatives as ameliorating agents for digestive tract movements. US 5466715 and US 5258407 describe the preparation of 3,4-disubstituted phenol immunostimulants. JP 58069812 describes hypoglycemic
25 pharmaceuticals containing benzamide derivatives. US 3950351 describes 2-benzamido-5-nitrothiazoles and Cavier *et al* [Eur J Med Chem - Chim Ther (1978) 13(6), 539-43] discuss the biological interest of these compounds.

Pending International application Number: PCT/GB02/02873 describes a group of benzoyl amino pyridyl carboxylic acids which are activators of the enzyme glucokinase
30 (GLK). We have surprisingly found a small selection of these compounds which have a superior level of drug in plasma following oral administration which is due to improved aqueous solubility and decreased levels of plasma binding, whilst retaining high potency for

the GLK enzyme. This makes this sub-group of compounds particularly suitable for use in the treatment or prevention of a disease or medical condition mediated through GLK.

Thus, according to the first aspect of the invention there is provided a compound of Formula (I):




Formula (I)

wherein:

R^1 is selected from hydrogen and C_{1-4} alkyl;

R^2 is selected from: $R^4-C(R^{5a}R^{5b})-$, $R^4=C(R^6)-$ and $R^{7a}C(R^{7b})=C(R^6)-$;

10 R^3-X- is selected from: methyl, methoxymethyl and



R^4 is selected from C_{1-4} alkyl, phenyl, C_{3-6} cycloalkyl and heteroaryl, wherein R^4 is optionally substituted by one or two substituents independently selected from R^8 ;

R^{5a} and R^{5b} are independently selected from hydrogen, fluoro and C_{1-4} alkyl;

15 R^6 is selected from hydrogen and C_{1-4} alkyl;

R^{7a} and R^{7b} are independently selected from C_{1-4} alkyl wherein R^{7a} and R^{7b} are optionally substituted by one or two substituents independently selected from R^8 ;

R^8 is selected from C_{1-3} alkyl, C_{1-3} alkoxy, fluoro and chloro;

with the proviso that:

20 (i) at least one of R^{5a} and R^{5b} is fluoro; and

(ii) when R^2 is $R^4=C(R^6)-$ then R^4 is C_{3-6} cycloalkyl;

or a salt, pro-drug or solvate thereof.


Compounds of Formula (I) may form salts which are within the ambit of the invention.


Pharmaceutically acceptable salts are preferred although other salts may be useful in, for

25 example, isolating or purifying compounds.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups. For example, " C_{1-4} alkyl" includes propyl, isopropyl and *t*-butyl. For the avoidance of doubt, an alkyl chain can be joined to the rest of the molecule at the end of the alkyl

chain or in the middle of an alkyl chain, i.e. the definition of "alkyl" includes the following structures:



wherein  represents the point of attachment to the rest of the molecule.

5 The term "**heteroaryl**" refers to is an unsaturated, monocyclic ring containing 5-6 carbon atoms of which at least one atom is replaced with nitrogen, sulphur or oxygen, wherein a sulphur atoms in a heterocyclic ring may be oxidised to S(O) or S(O)₂. A heteroaryl ring may, unless otherwise specified, be carbon or nitrogen linked, unless linking via nitrogen leads to a charged quaternary nitrogen.

10 Examples of heteroaryl include: thienyl, furanyl, thiazolyl, thiadiazolyl, triazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl. Preferred heteroaryl rings include: thienyl.

The term "**C₃₋₆cycloalkyl**" refers to a saturated carbocyclic ring containing between 3 to 6 carbon atoms, preferably between 5 and 6 carbon atoms. Examples of C₃₋₆cycloalkyl
15 include cyclohexyl, cyclopentyl, cyclobutyl or cyclopropyl. Preferably cyclopentyl or cyclohexyl. Most preferably cyclopentyl.

Examples of **C₁₋₄alkyl** include methyl, ethyl, propyl, isopropyl, *sec*-butyl and *tert*-butyl; examples of **C₁₋₃alkoxy** include methoxy, ethoxy, propoxy, and isopropoxy


It is to be understood that, insofar as certain of the compounds of Formula (I) defined
20 above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the property of stimulating GLK directly or inhibiting the GLK/GLKRP interaction. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active
25 starting materials or by resolution of a racemic form. It is also to be understood that certain compounds may exist in tautomeric forms and that the invention also relates to any and all tautomeric forms of the compounds of the invention which activate GLK.

Preferred compounds of Formula (I) are those wherein any one or more of the following
30 apply:

- (1) R^2 is $R^4-C(R^{5a}R^{5b})-$
- (2) R^2 is $R^4-C(R^{5a}R^{5b})-$ and R^4 is phenyl;
- (3) R^2 is $R^4-C(R^{5a}R^{5b})-$ and R^4 is heteroaryl;
- (4) R^2 is $R^4-C(R^{5a}R^{5b})-$ and R^4 is C_{3-6} cycloalkyl;
- 5 (5) R^2 is $R^4-C(R^{5a}R^{5b})-$ and both R^{5a} and R^{5b} are fluoro;
- (6) R^2 is $R^4=C(R^6)-$;
- (7) R^2 is $R^4=C(R^6)-$ and R^3-X- is methyl;
- (8) R^2 is $R^4=C(R^6)-$ and R^3-X- is methoxymethyl;
- (9) R^2 is $R^4-C(R^{5a}R^{5b})-$ and R^3-X- is methyl;
- 10 (10) R^2 is $R^4-C(R^{5a}R^{5b})-$ and R^3-X- is methoxymethyl;
- (11) R^4 is unsubstituted;
- (12) R^3-X- is methyl;
- (13) R^3-X- is methoxymethyl;
- (14) R^2 is $R^{7a}C(R^{7b})=C(R^6)-$.

15 According to a further feature of the invention there is provided the following preferred groups of compounds of the invention:

(I) a compound of Formula (Ia)




Formula (Ia)

20 wherein:

R^1 and R^2 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(II) a compound of Formula (Ib)




Formula (Ib)

25

wherein:

R^1 and R^2 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(III) a compound of Formula (Ic)




5

Formula (Ic)

wherein:

R^1 and R^2 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(IV) a compound of Formula (Id)




10

Formula (Id)

wherein:

R^1 and R^2 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

15 (V) a compound of Formula (Ie)



Formula (Ie)

wherein:

R^1 and R^2 , R^3 , R^4 , R^{5a} , and R^{5b} are as defined above in a compound of Formula (I);

20 or a salt, solvate or pro-drug thereof.

(VI) Compounds of Formula (I) wherein

R^1 is hydrogen;

- R^2 is selected from: $R^4-C(R^{5a}R^{5b})-$ and $R^4=C(R^6)-$;
 R^3-X- is selected from methyl and methoxymethyl;
 R^4 is selected from phenyl and C_{3-6} cycloalkyl, wherein R^4 is optionally substituted by one or two substituents independently selected from R^7 , preferably unsubstituted;
5 R^{5a} and R^{5b} are independently selected from hydrogen and fluoro;
 R^6 is hydrogen;
 R^7 is independently selected from C_{1-3} alkyl, C_{1-3} alkoxy, fluoro and chloro;
with the proviso that:
(i) at least one of R^{5a} and R^{5b} is fluoro, preferably both R^{5a} and R^{5b} are fluoro;
10 (ii) when R^2 is $R^4=C(R^6)-$ then R^4 is C_{3-6} cycloalkyl.

In a further aspect of the invention there is provided any one of the Examples, or a salt, solvate or pro-drug thereof. In a further aspect of the invention there is provided any two or more of the Examples, or a salt, solvate or pro-drug thereof.

- 15 Preferred compounds of the invention include any one, two or more of:
6- $\{[(3-[(2,2\text{-difluoro-2-phenylethyl)oxy}]-5-[(1S)\text{-1-methyl-2-(methyloxy)ethyl)oxy}]phenyl)carbonyl]amino\}$ pyridine-3-carboxylic acid
6- $\{[(3-[(2,2\text{-difluoro-2-phenylethyl)oxy}]-5-[(1\text{-methylethyl)oxy}]phenyl)carbonyl]amino\}$ pyridine-3-carboxylic acid
20 6- $\{[(3-[(2\text{-cyclopentylideneethyl)oxy}]-5-[(1S)\text{-1-methyl-2-(methyloxy)ethyl)oxy}]phenyl)carbonyl]amino\}$ pyridine-3-carboxylic acid
6- $\{[(3-[(2\text{-cyclopentylideneethyl)oxy}]-5-[(1\text{-methylethyl)oxy}]phenyl)carbonyl]amino\}$ pyridine-3-carboxylic acid

or a salt, solvate or pro-drug thereof.

- 25 The compounds of the invention may be administered in the form of a pro-drug. A pro-drug is a bioprecursor or pharmaceutically acceptable compound being degradable in the body to produce a compound of the invention (such as an ester or amide of a compound of the invention, particularly an *in vivo* hydrolysable ester). Various forms of prodrugs are known in the art. For examples of such prodrug derivatives, see:
30 a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen;

- c) H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- d) H. Bundgaard, *Advanced Drug Delivery Reviews*, 8, 1-38 (1992);
- e) H. Bundgaard, *et al.*, *Journal of Pharmaceutical Sciences*, 77, 285 (1988); and
- 5 f) N. Kakeya, *et al.*, *Chem Pharm Bull*, 32, 692 (1984).

The contents of the above cited documents are incorporated herein by reference.

Examples of pro-drugs are as follows. An in-vivo hydrolysable ester of a compound of the invention containing a carboxy or a hydroxy group is, for example, a pharmaceutically-acceptable ester which is hydrolysed in the human or animal body to produce the parent acid
10 or alcohol. Suitable pharmaceutically-acceptable esters for carboxy include C₁ to C₆alkoxymethyl esters for example methoxymethyl, C₁ to C₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃ to C₈cycloalkoxycarbonyloxy C₁ to C₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters, for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁-C₆alkoxycarbonyloxyethyl esters.

15 An in-vivo hydrolysable ester of a compound of the invention containing a hydroxy group includes inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the in-vivo hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy.

20 A selection of in-vivo hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl.

A suitable pharmaceutically-acceptable salt of a compound of the invention is, for
25 example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically-acceptable salt of a benzoxazinone derivative of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or
30 potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable

cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

A further feature of the invention is a pharmaceutical composition comprising a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) as defined above, or a salt, solvate or
5 prodrug thereof, together with a pharmaceutically-acceptable diluent or carrier.

According to another aspect of the invention there is provided a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) as defined above for use as a medicament.

Further according to the invention there is provided a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) for use in the preparation of a medicament for treatment of a disease
10 mediated through GLK, in particular type 2 diabetes.

The compound is suitably formulated as a pharmaceutical composition for use in this way.

According to another aspect of the present invention there is provided a method of treating GLK mediated diseases, especially diabetes, by administering an effective amount of
15 a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie), or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

Specific disease which may be treated by the compound or composition of the invention include: blood glucose lowering in Diabetes Mellitus type 2 without a serious risk of hypoglycaemia (and potential to treat type 1), dyslipidemia, obesity, insulin resistance,
20 metabolic syndrome X, impaired glucose tolerance.

As discussed above, thus the GLK/GLKRP system can be described as a potential "Diabesity" target (of benefit in both Diabetes and Obesity). Thus, according to another aspect of the invention there is provided the use of a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie), or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in
25 the combined treatment or prevention of diabetes and obesity.

According to another aspect of the invention there is provided the use of a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie), or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the treatment or prevention of obesity.

According to a further aspect of the invention there is provided a method for the
30 combined treatment of obesity and diabetes by administering an effective amount of a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie), or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

According to a further aspect of the invention there is provided a method for the treatment of obesity by administering an effective amount of a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie), or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

5 The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for
10 example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended
15 for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding
20 agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using
25 conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

30 Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation

products of an alkylene oxide with fatty acids (for example polyoxyethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or
5 condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions
10 may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid
15 paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by
20 the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of
25 oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan
30 monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous
5 or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

10 Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

15 For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the
20 particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information
25 on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) will naturally vary according to the nature and severity
30 of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.

In using a compound of the Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for

example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose
5 in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred.

The elevation of GLK activity described herein may be applied as a sole therapy or may involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous,
10 sequential or separate administration of the individual components of the treatment.

Simultaneous treatment may be in a single tablet or in separate tablets. For example in the treatment of diabetes mellitus chemotherapy may include the following main categories of treatment:

- 1) Insulin and insulin analogues;
- 15 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide), prandial glucose regulators (for example repaglinide, nateglinide);
- 3) Agents that improve incretin action (for example dipeptidyl peptidase IV inhibitors, and GLP-1 agonists);
- 4) Insulin sensitising agents including PPARgamma agonists (for example pioglitazone
20 and rosiglitazone), and agents with combined PPARalpha and gamma activity;
- 5) Agents that modulate hepatic glucose balance (for example metformin, fructose 1, 6 bisphosphatase inhibitors, glycogen phosphorylase inhibitors, glycogen synthase kinase inhibitors);
- 6) Agents designed to reduce the absorption of glucose from the intestine (for example
25 acarbose);
- 7) Agents that prevent the reabsorption of glucose by the kidney (SGLT inhibitors);
- 8) Agents designed to treat the complications of prolonged hyperglycaemia (for example aldose reductase inhibitors);
- 9) Anti-obesity agents (for example sibutramine and orlistat);
- 30 10) Anti-dyslipidaemia agents such as, HMG-CoA reductase inhibitors (eg statins); PPARα agonists (fibrates, eg gemfibrozil); bile acid sequestrants (cholestyramine);

cholesterol absorption inhibitors (plant stanols, synthetic inhibitors); bile acid absorption inhibitors (IBATi) and nicotinic acid and analogues (niacin and slow release formulations);


- 11) Antihypertensive agents such as, β blockers (eg atenolol, inderal); ACE inhibitors (eg lisinopril); Calcium antagonists (eg. nifedipine); Angiotensin receptor antagonists (eg
5 candesartan), α antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 12) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors; antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low molecular weight analogues, hirudin) and warfarin;
- 10 13) Agents which antagonise the actions of glucagon; and
- 14) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin) and steroidal anti-inflammatory agents (eg. cortisone).

According to another aspect of the present invention there is provided individual compounds produced as end products in the Examples set out below and salts, solvates and
15 pro-drugs thereof.


A compound of the invention, or a salt thereof, may be prepared by any process known to be applicable to the preparation of such compounds or structurally related compounds. Functional groups may be protected and deprotected using conventional methods. For examples of protecting groups such as amino and carboxylic acid protecting groups (as well as
20 means of formation and eventual deprotection), see T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis", Second Edition, John Wiley & Sons, New York, 1991.

Processes for the synthesis of compounds of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) are provided as a further feature of the invention. Thus, according to a further aspect of the
25 invention there is provided a process for the preparation of a compound of Formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) which comprises:

(a) reaction of an acid of Formula (IIIa) or activated derivative thereof with a compound of Formula (IIIb),



Formula (IIIa)




Formula (IIIb);

5 wherein P^1 is hydrogen or a protecting group such as C_{1-4} alkyl, (preferably methyl or ethyl);

or


(b) de-protection of a compound of Formula (IIIc),



Formula (IIIc)

10 wherein P^2 is a protecting group; or

(c) reaction of a compound of Formula (IIIId) with a compound of Formula (IIIe),




Formula (IIIId)

Formula (IIIe)

15 wherein X^1 is a leaving group and X^2 is a hydroxyl group or X^1 is a hydroxyl group and X^2 is a leaving group and wherein P^1 is hydrogen or a protecting group; or


(d) reaction of a compound of Formula (III_f) with a compound of Formula (III_g)

Formula (III_f)Formula (III_g)

wherein X³ is a leaving group and X⁴ is a hydroxyl group or X³ is a hydroxyl group and

5 X⁴ is a leaving group wherein P¹ is hydrogen or a protecting group; or

(e) reaction of a compound of Formula (III_h) with a compound of Formula (III_i),

Formula (III_h)Formula (III_i);

wherein X⁵ is a leaving group and wherein P¹ is hydrogen or a protecting group;

10 and thereafter, if necessary:

- i) converting a compound of Formula (I) into another compound of Formula (I);
- ii) removing any protecting groups;
- iii) forming a salt, pro-drug or solvate thereof.

Suitable leaving groups for processes a) to e) are well known to the skilled person and
15 include for example activated hydroxy leaving groups (such as mesylate and tosylate groups) and halo leaving groups such as fluoro, chloro or bromo.

Compounds of formulae (III_a) to (III_i) are commercially available, or may be made by any convenient process known in the art and/or as illustrated in the Examples herein. In general it will be appreciated that any aryl-O or alkyl-O bond may be formed by nucleophilic substitution or
20 metal catalysed processes, optionally in the presence of a suitable base.

Specific reaction conditions for the above reactions are as follows, wherein when P¹ is a protecting group P¹ is preferably C₁₋₄alkyl, for example methyl or ethyl:

Process a) – coupling reactions of amino groups with carboxylic acids to form an amide are well known in the art. For example,

(i) using an appropriate coupling reaction, such as a carbodiimide coupling reaction performed with EDAC in the presence of DMAP in a suitable solvent such as DCM, chloroform or DMF at room temperature; or

(ii) reaction in which the carboxylic group is activated to an acid chloride by reaction with oxalyl chloride in the presence of a suitable solvent such as methylene chloride. The acid chloride can then be reacted with a compound of Formula IIIb in the presence of a base, such as triethylamine or pyridine, in a suitable solvent such as chloroform or DCM at a temperature between 0°C and room temperature.

Process b) – de-protection reactions are well known in the art. Examples of P¹ include C₁₋₆alkyl and benzyl. Wherein P¹ is an C₁₋₆alkyl, the reaction can be performed in the presence of sodium hydroxide in the suitable solvent such as THF/water.

Process c) – compounds of Formula (III_d) and (III_e) can be reacted together in a suitable solvent, such as DMF or THF, with a base such as sodium hydride or potassium *tert*-butoxide, at a temperature in the range 0 to 100°C, optionally using metal catalysis such as palladium(II)acetate, palladium on carbon, copper(II)acetate or copper(I)iodide; Alternatively, compounds of Formula (III_d) and (III_e) can be reacted together in a suitable solvent, such as THF or DCM, with a suitable phosphine such as triphenylphosphine, and azodicarboxylate such as diethylazodicarboxylate;

Process d) – compounds of Formula (III_d) and (III_e) can be reacted together in a suitable solvent, such as DMF or THF, with a base such as sodium hydride or potassium *tert*-butoxide, at a temperature in the range 0 to 100°C, optionally using metal catalysis such as palladium(II)acetate, palladium on carbon, copper(II)acetate or copper(I)iodide, alternatively, compounds of Formula (III_d) and (III_e) can be reacted together in a suitable solvent, such as THF or DCM, with a suitable phosphine such as triphenylphosphine, and azodicarboxylate such as diethylazodicarboxylate;

Process e) – reaction of a compound of Formula (III_h) with a compound of Formula (III_i) can be performed in a polar solvent, such as DMF or a non-polar solvent such as THF with a strong base, such as sodium hydride or potassium *tert*-butoxide at a temperature between 0 and 100°C, optionally using metal catalysis, such as palladium(II)acetate, palladium on carbon, copper(II)acetate or copper(I)iodide.

During the preparation process, it may be advantageous to use a protecting group for a functional group within the molecule. Protecting groups may be removed by any convenient

method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

5 Specific examples of protecting groups are given below for the sake of convenience, in which "lower" signifies that the group to which it is applied preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically
10 mentioned is of course within the scope of the invention.

 A carboxy protecting group may be the residue of an ester-forming aliphatic or araliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms). Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (e.g. isopropyl, *t*-butyl); lower alkoxy lower alkyl groups
15 (e.g. methoxymethyl, ethoxymethyl, isobutoxymethyl; lower aliphatic acyloxy lower alkyl groups, (e.g. acetoxymethyl, propionyloxymethyl, butyryloxymethyl, pivaloyloxymethyl); lower alkoxycarbonyloxy lower alkyl groups (e.g. 1-methoxycarbonyloxyethyl, 1-ethoxycarbonyloxyethyl); aryl lower alkyl groups (e.g. *p*-methoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (e.g. trimethylsilyl and
20 *t*-butyldimethylsilyl); tri(lower alkyl)silyl lower alkyl groups (e.g. trimethylsilylethyl); and (2-6C)alkenyl groups (e.g. allyl and vinyloethyl).

 Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, metal- or enzymically-catalysed hydrolysis.

 Examples of hydroxy protecting groups include lower alkenyl groups (e.g. allyl); lower
25 alkanoyl groups (e.g. acetyl); lower alkoxycarbonyl groups (e.g. *t*-butoxycarbonyl); lower alkenyloxycarbonyl groups (e.g. allyloxycarbonyl); aryl lower alkoxycarbonyl groups (e.g. benzoyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl); tri lower alkyl/arylsilyl groups (e.g. trimethylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl); aryl lower alkyl groups (e.g. benzyl) groups; and
30 triaryl lower alkyl groups (e.g. triphenylmethyl).

 Examples of amino protecting groups include formyl, aralkyl groups (e.g. benzyl and substituted benzyl, e.g. *p*-methoxybenzyl, nitrobenzyl and 2,4-dimethoxybenzyl, and triphenylmethyl); di-*p*-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (e.g.

t-butoxycarbonyl); lower alkenyloxycarbonyl (e.g. allyloxycarbonyl); aryl lower alkoxy carbonyl groups (e.g. benzyloxycarbonyl, p-methoxybenzyloxycarbonyl, o-nitrobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl; trialkylsilyl (e.g. trimethylsilyl and t-butyl dimethylsilyl); alkylidene (e.g. methylidene); benzylidene and substituted benzylidene groups.

Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base, metal- or enzymically-catalysed hydrolysis, or photolytically for groups such as o-nitrobenzyloxycarbonyl, or with fluoride ions for silyl groups.

Examples of protecting groups for amide groups include aralkoxymethyl (e.g. benzyloxymethyl and substituted benzyloxymethyl); alkoxy methyl (e.g. methoxymethyl and trimethylsilylethoxymethyl); tri alkyl/arylsilyl (e.g. trimethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl); tri alkyl/arylsilyloxymethyl (e.g. t-butyl dimethylsilyloxymethyl, t-butyl diphenylsilyloxymethyl); 4-alkoxyphenyl (e.g. 4-methoxyphenyl); 2,4-di(alkoxy)phenyl (e.g. 2,4-dimethoxyphenyl); 4-alkoxybenzyl (e.g. 4-methoxybenzyl); 2,4-di(alkoxy)benzyl (e.g. 2,4-di(methoxy)benzyl); and alk-1-enyl (e.g. allyl, but-1-enyl and substituted vinyl e.g. 2-phenylvinyl).

Aralkoxymethyl, groups may be introduced onto the amide group by reacting the latter group with the appropriate aralkoxymethyl chloride, and removed by catalytic hydrogenation. Alkoxy methyl, tri alkyl/arylsilyl and tri alkyl/silyloxymethyl groups may be introduced by reacting the amide with the appropriate chloride and removing with acid; or in the case of the silyl containing groups, fluoride ions. The alkoxyphenyl and alkoxybenzyl groups are conveniently introduced by arylation or alkylation with an appropriate halide and removed by oxidation with ceric ammonium nitrate. Finally alk-1-enyl groups may be introduced by reacting the amide with the appropriate aldehyde and removed with acid.

The following examples are for illustration purposes and are not intended to limit the scope of this application. Each exemplified compound represents a particular and independent aspect of the invention. In the following non-limiting Examples, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at room temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon or nitrogen;

(iii) yields are given for illustration only and are not necessarily the maximum attainable;

(iv) the structures of the end-products of the Formula (I) were confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic
5 resonance chemical shift values were measured on the delta scale and peak multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad; q, quartet, quin, quintet;

(v) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), high-performance liquid chromatography (HPLC), infra-red
10 (IR) or NMR analysis;

(vi) Isolute silica cartridges refer to pre-packed silica cartridges (from 1g up to 70g) from IST (International Sorbent Technology), Hengoed, Mid Glamorgan, Wales UK, CF82 7RJ, eluted using a Flashmaster 2 system; Argonaut Technologies, Inc., Hengoed, Mid Glamorgan, Wales UK CF82 8AU;

15 (vii) Biotage cartridges refer to pre-packed silica cartridges (from 40g up to 400g), eluted using a biotage pump and fraction collector system; Biotage UK Ltd, Hertford, Herts, UK.


(viii) Celite refers to diatomaceous earth.

20 Abbreviations

DCM	dichloromethane;
DEAD	diethyldiazocarbonylate;
DIAD	di-i-propyl azodicarbonylate;
DIPEA	di-isopropylethylamine
25 DMSO	dimethyl sulphoxide;
DMF	dimethylformamide;
EDAC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride;
LCMS	liquid chromatography / mass spectroscopy;
30 RT	room temperature; and
THF	tetrahydrofuran.

EXAMPLE 1

6-{[(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1*S*)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl]amino}pyridine-3-carboxylic acid



5


To a solution of methyl 6-{[(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1*S*)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl]amino}pyridine-3-carboxylate (375mg, 0.75 mmol) in THF (5ml) was added distilled water (1.9ml) and sodium hydroxide solution (1.9ml of 1M, 1.9 mmol, ~2.5 eq). Methanol (2 drops) was added to aid solubility, and the mixture stirred at ambient temperature for 2 hours. The reaction mixture was neutralised with hydrochloric acid solution (1.9ml of 1M) and the THF partially removed in *vacuo*; more water was added, and the resulting solid was filtered off and washed with more distilled water. After partial drying, the solid was suspended in acetonitrile (4ml) and stirred gently for ~1hr; the solid was filtered, washed with more acetonitrile and dried to give 6-{[(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1*S*)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl]amino}pyridine-3-carboxylic acid as a colourless solid,

15

¹H NMR δ (d₆-DMSO): 1.2 (d, 3H), 3.25 (s, 3H), 3.4-3.55 (m, 2H), 4.6-4.8 (m, 3H), 6.75 (m, 1H), 7.25 (d, 2H), 7.55 (m, 3H), 7.65 (m, 2H), 8.3 (s, 2H), 8.85 (s, 1H), 11.05 (br s, 1H);
m/z 487 (M+H)⁺, 485 (M-H)⁻


20

Intermediates for the preparation of Example 1 were prepared according to the following scheme:



as described below.

Methyl 6-([(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1S)-1-methyl-2-(methyloxy)ethyl]oxy)phenyl]carbonyl)amino}pyridine-3-carboxylate



A solution of methyl 6-([(3-hydroxy-5-[(1S)-1-methyl-2-(methyloxy)ethyl]oxy)phenyl]carbonyl)amino}pyridine-3-carboxylate (360mg, 1mmol) in acetone (7ml) and DMF (2ml)
 10 was treated sequentially with potassium carbonate (414mg, 3mmol, 3 eq) and 2,2-difluoro-2-phenylethyl trifluoromethanesulfonate (432mg, 1.5mmol, 1.5 eq). The resulting suspension was stirred for 3 days at ambient temperature, adding extra sulfonate reagent (2x250mg portions on consecutive days).


The reaction mixture was then diluted with ethyl acetate and washed sequentially with water
 15 (twice) and brine, dried (MgSO₄) and evaporated to give the crude product (1g) as a brown oil. This was chromatographed (10g Isolute silica cartridge, eluting with hexane containing ethyl acetate, 15% increasing to 20%) to give methyl 6-([(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1S)-1-methyl-2-(methyloxy)ethyl]oxy)phenyl]carbonyl)amino}pyridine-3-carboxylate (390mg) as a colourless gum,

^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 3.25 (s, 3H), 3.4-3.55 (m, 2H), 3.85 (s, 3H), 4.6 – 4.8 (m, 3H), 6.8 (s, 1H), 7.25 (d, 2H), 7.55 (m, 3H), 7.65 (m, 2H), 8.35 (s, 2H), 8.9 (s, 1H), 11.1 (br s, 1H);

m/z 499 (M+H) $^+$, 501 (M-H) $^-$

5

Methyl 6-[(3-hydroxy-5-[(1*S*)-1-methyl-2-(methoxy)ethyl]oxy]phenyl)carbonyl]amino]pyridine-3-carboxylate




To a stirred solution of methyl 6-[(3-[(1*S*)-1-methyl-2-(methoxy)ethyl]oxy)-5-
 10 [(phenylmethyl)oxy]phenyl)carbonyl]amino]pyridine-3-carboxylate (0.038 mol) in THF (85 mL) was added methanol (85 mL). Palladium on charcoal catalyst (1.7g of 10% w/w) was added under an argon atmosphere, and the resulting suspension stirred at ambient temperature overnight in an atmosphere of hydrogen. The catalyst was filtered off through celite, washed with THF, and the filtrate evaporated to give a pale brown solid. This was triturated with ether
 15 to give the desired compound (72% yield).

^1H NMR δ (d_6 -DMSO): 1.25 (d, 3H), 3.3 (s, 3H), 3.45 (m, 2H), 3.85 (s, 3H), 4.65 (m, 1H), 6.55 (m, 1H), 6.95 (m, 1H), 7.1 (m, 1H), 8.3 (m, 2H), 8.9 (m, 1H), 11.0, (s, 1H).

m/z 361 (M+H) $^+$, 359 (M-H) $^-$

20 Methyl 6-[(3-[(1*S*)-1-methyl-2-(methoxy)ethyl]oxy)-5-[(phenylmethyl)oxy]phenyl)carbonyl]amino]pyridine-3-carboxylate




To a stirred solution of 3-[(1*S*)-1-methyl-2-(methoxy)ethyl]oxy)-5-[(phenylmethyl)oxy]
 benzoic acid (75.9 mmol) in DCM (250 mL) containing DMF (1 mL), oxalyl chloride was
 25 added dropwise under argon (151.7 mmol), and the resulting solution stirred for 4 hours. The solution was then evaporated *in vacuo*, azeotroped with more DCM (3x100 mL), and the residue dried under high vacuum to give the acid chloride, which was used without characterisation.

The acid chloride from above (approx. 75.9 mmol) was dissolved in THF (100 mL) and added
 30 under argon to a stirred solution of methyl 6-aminonicotinate (91.1 mmol) in a mixture of


THF (100 mL) and pyridine (100 mL). The reaction mixture was stirred overnight, and then most of the solvent removed *in vacuo*. The residue was taken up in ethyl acetate (250 mL), and the suspension washed sequentially with 1M citric acid (2 portions, until washings acidic) and brine; the resulting solution was dried (MgSO₄) and evaporated to give the crude product
 5 as a brown gum. This was chromatographed (400 g Biotage silica cartridge, eluting with hexane containing ethyl acetate, 20% v/v) to give the desired compound (50% yield).
¹H NMR δ (d₆-DMSO): 1.21 (d, 3H), 3.47 (m, 2H), 3.86 (s, 3H), 3.72 (m, 1H), 5.16 (s, 2H), 6.78 (t, 1H), 7.23 (s, 1H), 7.29 (s, 1H), 7.31-7.49 (m, 5H), 8.32 (s, 2H), 8.90 (app t, 1H), 11.15 (s, 1H).
 10 *m/z* 451.5 (M+H)⁺, 449.5 (M-H)⁻

3-[[*(1S)*-1-Methyl-2-(methyloxy)ethyl]oxy]-5-[(phenylmethyl)oxy]benzoic acid




A solution of methyl 3-[[*(1S)*-1-methyl-2-(methyloxy)ethyl]oxy]-5-[(phenylmethyl)oxy]
 15 benzoate (77.4 mmol) in a mixture of THF (232 mL) and methanol (232 mL) was treated with a solution of sodium hydroxide (2N) (232 mmol), and the reaction mixture stirred for 4 hours at ambient temperature. The resulting solution was diluted with water (250 mL) and most of the organic solvent removed *in vacuo*. The resulting suspension was washed with diethyl ether (3x200 mL) and the washings discarded. The resulting aqueous solution was acidified to pH 4
 20 with hydrochloric acid solution (2M) and extracted with ethyl acetate (2x200 mL); the extracts were combined, washed with brine, dried (MgSO₄) and evaporated to give the desired compound (99% yield).

¹H NMR δ (d₆-DMSO): 1.20 (d, 3H), 3.46 (m, 2H), 4.64 (m, 1H), 5.15 (s, 2H), 6.83 (app t, 1H), 7.06 (s, 1H), 7.13 (s, 1H), 7.30-7.49 (m, 5H), 12.67 (brs, 1H).

Methyl 3-[(1*S*)-1-methyl-2-(methyloxy)ethyl]oxy-5-[(phenylmethyl)oxy]benzoate

To a solution of methyl 3-hydroxy-5-[(phenylmethyl)oxy]benzoate (77.4 mmol) in THF was added polymer-supported triphenylphosphine (51.7g of 3 mmol/g loading, 155mmol) and (R)-
 5 (-)-1-methoxy-2-propanol (102 mmol). The stirred solution was blanketed with argon and cooled in an ice bath; a solution of diisopropyl azodicarboxylate (116 mmol) was added dropwise from a syringe over 10 minutes. After addition the solution was stirred for 20 minutes and then filtered, washing the residue with THF (500 mL); the filtrate and washings were combined and evaporated to give crude desired compound which was used in the next
 10 step without further purification.


¹H NMR δ (d₆-DMSO): 3.26 (s, 3H), 3.44 (m, 2H), 3.82 (s, 3H), 4.63 (m, 1H), 5.14 (s, 2H), 6.85 (s, 1H), 7.05 (s, 1H), 7.11 (s, 1H), 7.30-7.47 (m, 5H); the spectrum also contained signals consistent with a small amount of bis(1-methylethyl)hydrazine-1,2-dicarboxylate.

15 Methyl 3-hydroxy-5-[(phenylmethyl)oxy]benzoate

To a stirred solution of methyl 3,5-dihydroxybenzoate (5.95 mol) in DMF (6 L) was added potassium carbonate (9 mol), and the suspension stirred at ambient temperature under argon. To this was added benzyl bromide (8.42 mol) slowly over 1 hour, with a slight exotherm, and
 20 the reaction mixture stirred overnight at ambient temperature. It was then quenched cautiously with ammonium chloride solution (5 L) followed by water (35 L). The aqueous suspension was extracted with DCM (1x3 L and 2x5 L). The combined extracts were washed with water (10 L) and dried overnight (MgSO₄). The solution was evaporated in *vacuo*, and the crude product chromatographed in three batches (flash column, 3x2 kg silica, eluting with a gradient
 25 consisting of hexane containing 10% DCM, to neat DCM, to DCM containing 50% ethyl acetate) to eliminate starting material; the crude eluant was then chromatographed in 175 g batches (Amicon HPLC, 5 kg normal-phase silica, eluting with *iso*-hexane containing 20% v/v of ethyl acetate) to give the desired compound (21% yield).


^1H NMR δ (d_6 -DMSO): 3.8 (s, 3H), 5.1 (s, 2H), 6.65 (m, 1H), 7.0 (m, 1H), 7.05 (m, 1H), 7.3-7.5 (m, 5H), 9.85 (brs, 1H).

The requisite 2,2-difluoro-2-phenylethyl trifluoromethane sulfonate starting material was
5 prepared according to the following scheme:



as described below:

2,2-Difluoro-2-phenylethyl trifluoromethanesulfonate



10

To a cooled, stirred solution of 2,2-difluoro-2-phenylethanol (1.6g, 10mmol) and diisopropylethylamine (DIPEA, 2.1ml, 12mmol, 1.2 eq) in DCM (50ml) was added trifluoromethane sulfonic anhydride (2.0ml, 12 mmol, 1.2 eq), and the solution stirred for 2 hrs. Further DIPEA (0.5 ml, 3mmol) and triflic anhydride (0.5ml, 3mmol) were added, and the
15 reaction mixture stirred a further 2 hrs. It was then washed sequentially with water (twice) and brine, dried (MgSO_4) and evaporated to give the crude product as a dark brown oil; this was chromatographed (20g Isolute silica cartridge, eluting with hexane containing 5% v/v of ethyl acetate) to give 2,2-difluoro-2-phenylethyl trifluoromethanesulfonate as a pale brown oil which was used immediately without characterisation.


20 The requisite 2,2-difluoro-2-phenylethanol was prepared according to the method given in WO 98/20878, starting from methyl α,α difluorophenyl acetate (WJ Middleton et al, J. Org. Chem. (1980), 45, 2883-2887).

Using an analogous method to that described above, **Example 1.1** was also prepared:

No	Structure	MS	NMR
1.1		(M+H) ⁺ 457 (M-H) ⁻ 455	^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 4.6-4.8 (m, 3H), 6.75 (m, 1H), 7.2 (m, 1H), 7.25 (m, 1H), 7.55 (m, 3H), 7.65 (m, 2H), 8.3 (s, 2H), 8.9 (s, 1H), 11.05 (br s, 1H).

The appropriate intermediates for the preparation of Example 1.1 were prepared using an analogous method to those used for the preparation of intermediates for the preparation of Example 1, unless otherwise stated:


5 Methyl 6-[(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1-methylethyl)oxy]phenyl)carbonyl)amino]pyridine-3-carboxylate



^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 3.85 (s, 3H), 4.6 – 4.8 (m, 3H), 6.75 (m, 1H), 7.2 (m, 1H), 7.25 (m, 1H), 7.45-7.55 (m, 3H), 7.65 (m, 2H), 8.35 (m, 2H), 8.9 (s, 1H), 11.1 (br s, 1H); m/z 471 ($M+H$) $^+$

10


Methyl 6-[(3-hydroxy-5-[(1-methylethyl)oxy]phenyl)carbonyl)amino]pyridine-3-carboxylate



^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 3.85 (s, 3H), 4.65 (m, 1H), 6.55 (m, 1H), 6.95 (m, 1H), 7.1 (m, 1H), 8.3 (s, 2H), 8.9 (s, 1H), 9.7 (s, 1H), 11.0, (s, 1H).


15 m/z 331 ($M+H$) $^+$, 329 ($M-H$) $^-$

Methyl 6-[(3-benzyloxy-5-[(1-methylethyl)oxy]phenyl)carbonyl)amino]pyridine-3-carboxylate




20 ^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 3.85 (s, 3H), 4.7 (m, 1H), 5.2 (s, 2H), 6.75 (m, 1H), 7.2 (m, 1H), 7.3-7.5 (m, 6H), 8.35 (s, 2H), 8.90 (s, 1H), 11.15 (br s, 1H)

3-[(1-methylethyl)oxy]-5-[(phenylmethyl)oxy]benzoic acid



^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 4.65 (m, 1H), 5.15 (s, 2H), 6.8 (m, 1H), 7.05 (m, 1H), 7.15 (m, 1H), 7.30-7.5 (m, 5H), 12.95 (s br, 1H)


Methyl 3-[(1-methylethyl)oxy]-5-[(phenylmethyl)oxy]benzoate



5 A solution of methyl 3-isopropoxy-5-hydroxy-benzoate (25.0g, 119mmol) in DMF (250ml) was treated with potassium carbonate (41.1g, 297 mmol, 2.5 eq) and benzyl bromide (17ml, 143 mmol, 1.2 eq), and the resulting suspension heated at 60°C for 5 hrs. The solvent was removed *in vacuo* and the residue suspended in water (200ml); this was extracted with ethyl acetate (2x250 ml). The combined extracts were washed sequentially with water (4x150ml) and brine (2x100ml), dried (MgSO_4) and evaporated to give methyl 3-isopropoxy-5-benzyloxy benzoate (37.5g) as a yellow oil which contained traces ethyl acetate, benzyl alcohol and benzyl bromide,

15 ^1H NMR δ (d_6 -DMSO): 1.2 (d, 6H), 3.85 (s, 3H), 4.65 (m, 1H), 5.15 (s, 2H), 6.85 (m, 1H), 7.05 (m, 1H), 7.15 (m, 1H), 7.3-7.5 (m, 5H).


Methyl 3-[(1-methylethyl)oxy]-5-hydroxy benzoate



20 ^1H NMR δ (d_6 -DMSO): 1.2 (d, 6H), 3.8 (s, 3H), 4.55 (m, 1H), 6.55 (m, 1H), 6.9 (m, 1H), 6.95 (m, 1H).

EXAMPLE 2

6-{[(3-[(2-cyclopentylideneethyl)oxy]-5-{[(1S)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl]amino}pyridine-3-carboxylic acid




25 Example 2 was prepared from the corresponding ester, methyl 6-{[(3-[(2-cyclopentylideneethyl)oxy]-5-{[(1S)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl] amino}pyridine-3-carboxylate using an analogous method to the preparation of Example 1

^1H NMR δ (d_6 -DMSO): 1.24 (s, 3H), 1.55-1.72 (m, 4H), 2.30 (app q, 4H), 3.30 (s, 3H obscured by solvent peak), 3.49 (qd, 2H), 4.57 (d, 2H), 4.75 (m, 1H), 5.55 (m, 1H), 6.70 (s, 1H), 7.18 (s, 1H), 7.22 (s, 1H), 8.31 (s, 2H), 8.90 (s, 1H), 11.09 (s, 1H), 13.17 (s br, 1H)
 m/z 441.5 ($\text{M}+\text{H}^+$), 439.5 ($\text{M}-\text{H}^-$)

5


Intermediates for the preparation of Example 2 were prepared according to the following scheme:



as described below:

10

Methyl 6-{[(3-[(2-cyclopentylidene ethyl)oxy]-5-{[(1S)-1-methyl-2-(methyloxy)ethyl]oxy} phenyl)carbonyl] amino }pyridine-3-carboxylate




To a stirred suspension of methyl 6-{[(3-hydroxy-5-{[(1S)-1-methyl-2-(methyloxy)ethyl]oxy} phenyl)carbonyl]amino }pyridine-3-carboxylate (900mg, 2.5 mmol), 2-cyclopentylidene ethanol (382 mg, 3.4 mmol, 1.4 eq) and polymer-supported triphenyl phosphine (approx. 3 mmol/g, 1.5 g, approx 3 eq) in DCM (30 ml), under argon, was added di-*tert*-butyl azodicarboxylate (DTAD, 1.29g, 5.6 mmol, 2.2 eq), and the reaction mixture stirred overnight at ambient temperature. The resin was removed by filtration and washed with ethyl acetate and THF; the filtrate and washings were combined and evaporated *in vacuo*, and the residue

20

trituated (ether/isohexane 1:1) to give methyl 6-{[(3-[(2-cyclopentylidene ethyl)oxy]-5-
 {[(1*S*)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl] amino}pyridine-3-carboxylate
¹H NMR δ (d₆-DMSO): 1.22 (d, 3H), 1.60 (m, 4H), 2.29 (d, 4H), 3.3 (s, 3H), 3.47 (m, 2H),
 3.45 (d, 2H), 3.88 (s, 3H), 4.71 (m, 1H), 5.53 (m, 1H), 6.68 (m, 1H), 7.18 (s, 1H), 7.20 (s,
 5 1H), 8.33 (s, 2H), 8.90 (s, 1H), 11.12 (br s, 1H)
 m/z 455.5 (M+H)⁺


Methyl 6-[(3-hydroxy-5-[(1*S*)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl)carbonyl]amino}
pyridine-3-carboxylate



10

This was prepared as described in the intermediates for Example 1 above.

2-Cyclopentylidene ethanol




15 This was prepared as described in International patent application number, WO 01/68603 on page 63.

Using an analogous method to that described above, **Example 2.1** was also prepared:

No	Structure	MS	NMR
2.1		(M+H) ⁺ 411	¹ H NMR δ (d ₆ -DMSO): 1.26 (d, 6H), 1.61 (m, 4H), 2.26 (d, 4H), 4.55 (d, 2H), 4.71 (septet, 1H), 5.53 (br m, 1H), 6.64 (t, 1H), 7.16 (m, 2H), 8.29 (s, 2H), 8.97 (s, 1H), 11.05 (br s, 1H), COOH not seen


20 The appropriate intermediates for the preparation of Example 2.1 were prepared using an analogous method to those used for the preparation of intermediates for the preparation of Example 2:

Methyl 6-([3-[(2-cyclopentylideneethyl)oxy]-5-[(1-methylethyl)oxy]phenyl]carbonyl)amino]pyridine-3-carboxylate



¹H NMR δ (d₆-DMSO): 1.27 (d, 6H), 1.60 (m, 4H), 2.28 (m, 4H), 3.86 (s, 3H), 4.54 (d, 2H),
 5 4.70 (septet, 1H), 5.52 (m, 1H), 6.63 (t, 1H), 7.15 (m, 2H), 8.32 (s, 2H), 8.88 (s, 1H), 11.10
 (br s, 1H)
 m/z 425 (M+H)⁺

Methyl 6-([3-hydroxy-5-[(1-methylethyl)oxy]phenyl]carbonyl)amino]pyridine-3-carboxylate



10

This was prepared as described in the intermediates for Example 1.1 above.

BIOLOGICAL

Tests:

15 The biological effects of the compounds of the invention may be tested in the following way:

(1) Enzymatic activity of GLK may be measured by incubating GLK, ATP and glucose. The rate of product formation may be determined by coupling the assay to a G-6-P dehydrogenase, NADP/NADPH system and measuring the linear increase in optical density at
 20 340nm (Matschinsky et al 1993). Activation of GLK by compounds can be assessed using this assay in the presence or absence of GLKRP as described in Brocklehurst et al (Diabetes 2004, 53, 535-541).

(2) A GLK/GLKRP binding assay for measuring the binding interactions between
 25 GLK and GLKRP (RP = regulatory protein). The method may be used to identify compounds which modulate GLK by modulating the interaction between GLK and GLKRP. GLKRP and GLK are incubated with an inhibitory concentration of F-6-P, optionally in the presence of test

compound, and the extent of interaction between GLK and GLKRP is measured. Compounds which either displace F-6-P or in some other way reduce the GLK/GLKRP interaction will be detected by a decrease in the amount of GLK/GLKRP complex formed. Compounds which promote F-6-P binding or in some other way enhance the GLK/GLKRP interaction will be
5 detected by an increase in the amount of GLK/GLKRP complex formed. A specific example of such a binding assay is described below

GLK/GLKRP scintillation proximity assay

Recombinant human GLK and GLKRP were used to develop a "mix and measure" 96
10 well SPA (scintillation proximity assay) as described in WO01/20327 (the contents of which are incorporated herein by reference). GLK (Biotinylated) and GLKRP are incubated with streptavidin linked SPA beads (Amersham) in the presence of an inhibitory concentration of radiolabelled [3H]F-6-P (Amersham Custom Synthesis TRQ8689), giving a signal. Compounds which either displace the F-6-P or in some other way disrupt the GLK / GLKRP
15 binding interaction will cause this signal to be lost.

Binding assays were performed at room temperature for 2 hours. The reaction mixtures contained 50mM Tris-HCl (pH 7.5), 2mM ATP, 5mM MgCl₂, 0.5mM DTT, recombinant biotinylated GLK (0.1 mg), recombinant GLKRP (0.1 mg), 0.05mCi [3H] F-6-P (Amersham) to give a final volume of 100ml. Following incubation, the extent of
20 GLK/GLKRP complex formation was determined by addition of 0.1mg/well avidin linked SPA beads (Amersham) and scintillation counting on a Packard TopCount NXT.

(3) A F-6-P / GLKRP binding assay for measuring the binding interaction between GLKRP and F-6-P. This method may be used to provide further information on the
25 mechanism of action of the compounds. Compounds identified in the GLK/GLKRP binding assay may modulate the interaction of GLK and GLKRP either by displacing F-6-P or by modifying the GLK/GLKRP interaction in some other way. For example, protein-protein interactions are generally known to occur by interactions through multiple binding sites. It is thus possible that a compound which modifies the interaction between GLK and GLKRP
30 could act by binding to one or more of several different binding sites.

The F-6-P / GLKRP binding assay identifies only those compounds which modulate the interaction of GLK and GLKRP by displacing F-6-P from its binding site on GLKRP.

GLKRP is incubated with test compound and an inhibitory concentration of F-6-P, in the absence of GLK, and the extent of interaction between F-6-P and GLKRP is measured. Compounds which displace the binding of F-6-P to GLKRP may be detected by a change in the amount of GLKRP/F-6-P complex formed. A specific example of such a binding assay is
5 described below

F-6-P / GLKRP scintillation proximity assay

Recombinant human GLKRP was used to develop a "mix and measure" 96 well scintillation proximity assay) as described in WO01/20327 (the contents of which are
10 incorporated herein by reference). FLAG-tagged GLKRP is incubated with protein A coated SPA beads (Amersham) and an anti-FLAG antibody in the presence of an inhibitory concentration of radiolabelled [3H]F-6-P. A signal is generated. Compounds which displace the F-6-P will cause this signal to be lost. A combination of this assay and the GLK/GLKRP binding assay will allow the observer to identify compounds which disrupt the GLK/GLKRP
15 binding interaction by displacing F-6-P.

Binding assays were performed at room temperature for 2 hours. The reaction mixtures contained 50mM Tris-HCl (pH 7.5), 2mM ATP, 5mM MgCl₂, 0.5mM DTT, recombinant FLAG tagged GLKRP (0.1 mg), Anti-Flag M2 Antibody (0.2mg) (IBI Kodak), 0.05mCi [3H] F-6-P (Amersham) to give a final volume of 100ml. Following incubation, the
20 extent of F-6-P/GLKRP complex formation was determined by addition of 0.1mg/well protein A linked SPA beads (Amersham) and scintillation counting on a Packard TopCount NXT.

Production of recombinant GLK and GLKRP:

25 *Preparation of mRNA*

Human liver total mRNA was prepared by polytron homogenisation in 4M guanidine isothiocyanate, 2.5mM citrate, 0.5% Sarkosyl, 100mM b-mercaptoethanol, followed by centrifugation through 5.7M CsCl, 25mM sodium acetate at 135,000g (max) as described in Sambrook J, Fritsch EF & Maniatis T, 1989.

30 Poly A⁺ mRNA was prepared directly using a FastTrack™ mRNA isolation kit (Invitrogen).

PCR amplification of GLK and GLKRP cDNA sequences

Human GLK and GLKRP cDNA was obtained by PCR from human hepatic mRNA using established techniques described in Sambrook, Fritsch & Maniatis, 1989. PCR primers were designed according to the GLK and GLKRP cDNA sequences shown in Tanizawa et al
5 1991 and Bonthron, D.T. *et al* 1994 (later corrected in Warner, J.P. 1995).

Cloning in Bluescript II vectors

GLK and GLKRP cDNA was cloned in E. coli using pBluescript II, (Short et al 1998) a recombinant cloning vector system similar to that employed by Yanisch-Perron C *et al*
10 (1985), comprising a colEI-based replicon bearing a polylinker DNA fragment containing multiple unique restriction sites, flanked by bacteriophage T3 and T7 promoter sequences; a filamentous phage origin of replication and an ampicillin drug resistance marker gene.

Transformations

15 E. Coli transformations were generally carried out by electroporation. 400 ml cultures of strains DH5a or BL21(DE3) were grown in L-broth to an OD 600 of 0.5 and harvested by centrifugation at 2,000g. The cells were washed twice in ice-cold deionised water, resuspended in 1ml 10% glycerol and stored in aliquots at -70°C. Ligation mixes were desalted using Millipore V series™ membranes (0.0025mm pore size). 40ml of cells were
20 incubated with 1ml of ligation mix or plasmid DNA on ice for 10 minutes in 0.2cm electroporation cuvettes, and then pulsed using a Gene Pulser™ apparatus (BioRad) at 0.5kVcm⁻¹, 250mF, 250 ?. Transformants were selected on L-agar supplemented with tetracycline at 10mg/ml or ampicillin at 100mg/ml.

25 Expression

GLK was expressed from the vector pTB375NBSE in E.coli BL21 cells,, producing a recombinant protein containing a 6-His tag immediately adjacent to the N-terminal methionine. Alternatively, another suitable vector is pET21(+)DNA, Novagen, Cat number 697703. The 6-His tag was used to allow purification of the recombinant protein on a column
30 packed with nickel-nitrilotriacetic acid agarose purchased from Qiagen (cat no 30250).

GLKRP was expressed from the vector pFLAG CTC (IBI Kodak) in E.coli BL21 cells, producing a recombinant protein containing a C-terminal FLAG tag. The protein was purified

initially by DEAE Sepharose ion exchange followed by utilisation of the FLAG tag for final purification on an M2 anti-FLAG immunoaffinity column purchased from Sigma-Aldrich (cat no. A1205).

5 Biotinylation of GLK:

GLK was biotinylated by reaction with biotinamidocaproate N-hydroxysuccinimide ester (biotin-NHS) purchased from Sigma-Aldrich (cat no. B2643). Briefly, free amino groups of the target protein (GLK) are reacted with biotin-NHS at a defined molar ratio forming stable amide bonds resulting in a product containing covalently bound biotin. Excess, non-
10 conjugated biotin-NHS is removed from the product by dialysis. Specifically, 7.5mg of GLK was added to 0.31mg of biotin-NHS in 4mL of 25mM HEPES pH7.3, 0.15M KCl, 1mM dithiothreitol, 1mM EDTA, 1mM MgCl₂ (buffer A). This reaction mixture was dialysed against 100mL of buffer A containing a further 22mg of biotin-NHS. After 4hours excess biotin-NHS was removed by extensive dialysis against buffer A.

15

Measurement of plasma levels and plasma protein binding following oral administration to rats

Administration of compounds to rats and sampling of plasma

Planetary Milled compounds [15mins, 500rpm, 5 Zirconium Balls, in a Pulverisette 7
20 Mill (Glen Creston Ltd, Stanmore, Middlesex, UK)] were suspended in 0.5% HPMC Tween and dosed to High Fat Fed (Research Diets, D12451, ad lib feeding 14 days) Female Alderley Park Zucker or Alderley Park Wistar rats at rate of 5mls/kg, at doses between 0.3 and 10mg/kg by oral gavage.

Samples of plasma were obtained either by conscious blood sampling or terminal blood
25 sampling as follows:

Conscious blood sampling (for compound level or blood chemistry) - Intravenous blood samples were taken from tail vein using 600µl Starstedt Multivette (EDTA) and 22G needle at the required time point. Samples were kept on ice and centrifuged at 3000rpm for 10 minutes within 15-30 minutes of withdrawal. The plasma was aspirated and stored at -20°C

30 Terminal blood sampling for compound level or blood chemistry - At the end of experiment animals were euthanased by exposure to CO₂/O₂. Blood sample were taken by

cardiac puncture. Samples were kept on ice and centrifuged at 3000rpm for 10 minutes within 15-30 minutes of withdrawal. The plasma was aspirated and stored at -20°C

Measurement of compound levels in rat plasma

5 25µl of rat plasma was added to wells in a 96 well protein precipitation plate (Varian inc. Palo Alto, California, USA). To each well was added 500µl of acetonitrile, containing 1ug/ml of (3-isopropoxy-5-benzyoxy-benzoyl)amino pyridine 3-carboxylic acid to act as an internal standard, to precipitate the plasma proteins. Then the plasma/solvent mixture was pulled through the precipitation plate under vacuum and the eluent was collected. The eluent
10 was evaporated to dryness using a centrifugal evaporator and reconstituted in 200µl of methanol:water:formic acid (60:40:0.1).

The reconstituted samples were then analysed using high performance liquid chromatography with tandem mass spectrometry detection (HPLC-MS-MS)". HPLC was performed using a Phenomenex Prodigy C8, 50x4.6, 5µm.column (Phenomenex,
15 Macclesfield, UK) at a flow rate of 1ml/minute using an injection volume of 10µl using the following gradient elution profile:

Mobile phase A	0.1% formic acid in water
Mobile phase B	0.1% formic acid in methanol
Mobile phase gradient	0 min 50% A
20	0.5 min 5% A
	2.5 min 5% A
	2.6 min 50% A
	3.0 min 50% A.

Mass spectroscopy was performed using an Applied Biosystems API3000 Mass
25 spectrometer (Applied Biosystems, Foster City, California, USA). Prior to the running of samples the mass spectrometer was optimised for the structure of the test compound.

The concentration of test samples was determined from the ratio of the peak height of the test sample to the peak height of the internal standard. The concentration of the test sample was calculated with reference to a standard curve relating the ratio to the concentration
30 prepared by using known concentrations of test sample added to samples of rat plasma using (3-isopropoxy-5-benzyoxy-benzoyl)amino pyridine 3-carboxylic acid as an internal standard, treated as described above.

Measurements of plasma protein binding of compounds

The plasma protein binding of compounds was measured using the equilibrium dialysis technique (W. Lindner et al, J.Chromatography, 1996, 677, 1-28). Compound was dialysed at a concentration of 20 μ M for 18 hours at 37°C with plasma and isotonic phosphate buffer pH 7.4 (1ml of each in the dialysis cell). A Spectrum® 20-cell equilibrium dialyser was used together with Teflon, semi-micro dialysis cells and Spectra/Por®2 membrane discs with a molecular weight cut off 12-14000 Dalton, 47mm (supplied by PerBio Science UK Ltd, Tattenhall, Cheshire). Plasma and buffer samples are removed following dialysis and analysed using HPLCUV/MS (high performance liquid chromatography with UV and mass spec
10 detection) to give the % free level in plasma.

Compounds of the invention have activate glucokinase with and EC_{50} of less than about 200nM, with a percentage free in plasma of between about 0.05% and about 1% and a peak blood levels (including both bound and free) of between about 0.5 μ M and about 10 μ M for a
15 normalised dose of 1mg compound per kilogram of rat body weight.

For example, Example 2 has the following values:

EC_{50}	% free in plasma	Peak Blood levels
78nM	0.42%	2.2 μ M

REFERENCES


- 20 1 Printz, R. L., Magnuson, M. A. and Granner, D. K. (1993) Annual Review of Nutrition 13, 463-96
- 2 DeFronzo, R. A. (1988) Diabetes 37, 667-87
- 3 Froguel, P., Zouali, H., Vionnet, N., Velho, G., Vaxillaire, M., Sun, F., Lesage, S., Stoffel, M., Takeda, J. and Passa, P. (1993) New England Journal of Medicine 328, 697-
25 702
- 4 Bell, G. I., Pilkis, S. J., Weber, I. T. and Polonsky, K. S. (1996) Annual Review of Physiology 58, 171-86

- 5 Velho, G., Petersen, K. F., Perseghin, G., Hwang, J. H., Rothman, D. L., Pueyo, M. E.,
Cline, G. W., Froguel, P. and Shulman, G. I. (1996) *Journal of Clinical Investigation* **98**,
1755-61
- 6 Christesen, H. B., Jacobsen, B. B., Odili, S., Buettger, C., Cuesta-Munoz, A., Hansen,
5 T., Brusgaard, K., Massa, O., Magnuson, M. A., Shiota, C., Matschinsky, F. M. and
Barbetti, F. (2002) *Diabetes* **51**, 1240-6
- 6a Gloyn, A.L., Noordam, K., Willemsen, M.A.A.P., Ellard, S., Lam, W.W.K., Campbell,
I. W., Midgley, P., Shiota, C., Buettger, C., Magnuson, M.A., Matschinsky, F.M., and
Hattersley, A.T.; *Diabetes* **52**: 2433-2440
- 10 7 Glaser, B., Kesavan, P., Heyman, M., Davis, E., Cuesta, A., Buchs, A., Stanley, C. A.,
Thornton, P. S., Permutt, M. A., Matschinsky, F. M. and Herold, K. C. (1998) *New
England Journal of Medicine* **338**, 226-30
- 8 Caro, J. F., Triester, S., Patel, V. K., Tapscott, E. B., Frazier, N. L. and Dohm, G. L.
(1995) *Hormone & Metabolic Research* **27**, 19-22
- 15 9 Desai, U. J., Slosberg, E. D., Boettcher, B. R., Caplan, S. L., Fanelli, B., Stephan, Z.,
Gunther, V. J., Kaleko, M. and Connelly, S. (2001) *Diabetes* **50**, 2287-95
- 10 Shiota, M., Postic, C., Fujimoto, Y., Jetton, T. L., Dixon, K., Pan, D., Grimsby, J.,
Grippe, J. F., Magnuson, M. A. and Cherrington, A. D. (2001) *Diabetes* **50**, 622-9
- 11 Ferre, T., Pujol, A., Riu, E., Bosch, F. and Valera, A. (1996) *Proceedings of the*
20 *National Academy of Sciences of the United States of America* **93**, 7225-30
- 12 Seoane, J., Barbera, A., Telemaque-Potts, S., Newgard, C. B. and Guinovart, J. J. (1999)
Journal of Biological Chemistry **274**, 31833-8
- 13 Moore, M. C., Davis, S. N., Mann, S. L. and Cherrington, A. D. (2001) *Diabetes Care*
24, 1882-7
- 25 14 Alvarez, E., Roncero, I., Chowen, J. A., Vazquez, P. and Blazquez, E. (2002) *Journal of
Neurochemistry* **80**, 45-53
- 15 Lynch, R. M., Tompkins, L. S., Brooks, H. L., Dunn-Meynell, A. A. and Levin, B. E.
(2000) *Diabetes* **49**, 693-700
- 16 Roncero, I., Alvarez, E., Vazquez, P. and Blazquez, E. (2000) *Journal of*
30 *Neurochemistry* **74**, 1848-57
- 17 Yang, X. J., Kow, L. M., Funabashi, T. and Mobbs, C. V. (1999) *Diabetes* **48**, 1763-
1772

- 18 Schuit, F. C., Huypens, P., Heimberg, H. and Pipeleers, D. G. (2001) *Diabetes* **50**, 1-11
- 19 Levin, B. E. (2001) *International Journal of Obesity* **25**, supplement 5, S68-S72.
- 20 Alvarez, E., Roncero, I., Chowen, J. A., Thorens, B. and Blazquez, E. (1996) *Journal of Neurochemistry* **66**, 920-7
- 5 21 Mobbs, C. V., Kow, L. M. and Yang, X. J. (2001) *American Journal of Physiology - Endocrinology & Metabolism* **281**, E649-54
- 22 Levin, B. E., Dunn-Meynell, A. A. and Routh, V. H. (1999) *American Journal of Physiology* **276**, R1223-31
- 23 Spanswick, D., Smith, M. A., Groppi, V. E., Logan, S. D. and Ashford, M. L. (1997)
- 10 24 Spanswick, D., Smith, M. A., Mirshamsi, S., Routh, V. H. and Ashford, M. L. (2000) *Nature* **390**, 521-5
- 25 Spanswick, D., Smith, M. A., Mirshamsi, S., Routh, V. H. and Ashford, M. L. (2000) *Nature Neuroscience* **3**, 757-8
- 26 Levin, B. E. and Dunn-Meynell, A. A. (1997) *Brain Research* **776**, 146-53
- 27 Levin, B. E., Govek, E. K. and Dunn-Meynell, A. A. (1998) *Brain Research* **808**, 317-9
- 15 28 Levin, B. E., Brown, K. L. and Dunn-Meynell, A. A. (1996) *Brain Research* **739**, 293-300
- 29 Rowe, I. C., Boden, P. R. and Ashford, M. L. (1996) *Journal of Physiology* **497**, 365-77
- 30 Fujimoto, K., Sakata, T., Arase, K., Kurata, K., Okabe, Y. and Shiraishi, T. (1985) *Life Sciences* **37**, 2475-82
- 20 31 Kurata, K., Fujimoto, K. and Sakata, T. (1989) *Metabolism: Clinical & Experimental* **38**, 46-51
- 32 Kurata, K., Fujimoto, K., Sakata, T., Etou, H. and Fukagawa, K. (1986) *Physiology & Behavior* **37**, 615-20

Claims:

1. A compound of Formula (I):



Formula (I)

wherein:

R^1 is selected from hydrogen and C_{1-4} alkyl;

R^2 is selected from: $R^4-C(R^{5a}R^{5b})-$, $R^4=C(R^6)-$ and $R^{7a}C(R^{7b})=C(R^6)-$;

R^3-X- is selected from methyl, methoxymethyl and

$H-C\equiv C-\frac{1}{2}$;

R^4 is selected from C_{1-4} alkyl, phenyl, C_{3-6} cycloalkyl and heteroaryl, wherein R^4 is optionally substituted by one or two substituents independently selected from R^8 ;

R^{5a} and R^{5b} are independently selected from hydrogen, fluoro and C_{1-4} alkyl;

R^6 is selected from hydrogen and C_{1-4} alkyl;

R^{7a} and R^{7b} are independently selected from C_{1-4} alkyl wherein R^{7a} and R^{7b} are

optionally substituted by one or two substituents independently selected from R^8 ;

R^8 is independently selected from C_{1-3} alkyl, C_{1-3} alkoxy, fluoro and chloro;


with the proviso that:

(i) at least one of R^{5a} and R^{5b} is fluoro; and

(ii) when R^2 is $R^4=C(R^6)-$ then R^4 is C_{3-6} cycloalkyl;

or a salt, pro-drug or solvate thereof.

2. A compound of formula (I) as claimed in Claim 1, which is a compound of Formula (Ia)




Formula (Ia)

wherein:

R^1 and R^2 are as in claim 1;

5 or a salt, solvate or pro-drug thereof.

3. A compound of formula (I) as claimed in Claim 1, which is a compound of a compound of Formula (Ic)



Formula (Ic)

wherein:

R^1 and R^2 are as defined in Claim 1;

or a salt, solvate or pro-drug thereof.

15 4. A compound as claimed in any of claims 1 to 3, wherein R^2 is $\text{R}^4\text{-C(R}^{5a}\text{R}^{5b})\text{-}$, or a salt, solvate or pro-drug thereof.

5. A compound as claimed in any of claims 1 to 3, wherein R^2 is $\text{R}^4\text{=C(R}^6\text{)-}$; or a salt, solvate or pro-drug thereof.

20

6. A compound of formula (I) as claimed in Claim 1, wherein

R^1 is hydrogen;

R^2 is selected from: $\text{R}^4\text{-C(R}^{5a}\text{R}^{5b})\text{-}$ and $\text{R}^4\text{=C(R}^6\text{)-}$;

$\text{R}^3\text{-X-}$ is selected from methyl and methoxymethyl;

25 R^4 is selected from phenyl and C_{3-6} cycloalkyl, wherein R^4 is optionally substituted by one or two substituents independently selected from R^7 ;

R^{5a} and R^{5b} are independently selected from hydrogen and fluoro;

R^6 is hydrogen;

R^7 is independently selected from C_{1-3} alkyl, C_{1-3} alkoxy, fluoro and chloro;

with the proviso that:

5 (iii) at least one of R^{5a} and R^{5b} is fluoro;

(iv) when R^2 is $R^4=C(R^6)-$ then R^4 is C_{3-6} cycloalkyl;

or a salt, solvate or pro-drug thereof.

7. A compound of formula (I) as claimed in Claim 6 wherein R^7 is unsubstituted; or a
10 salt, solvate or pro-drug thereof.

8. A compound of formula (I) as claimed in Claim 6 wherein both R^{5a} and R^{5b} are fluoro; or a salt, solvate or pro-drug thereof.

15 9. A compound of formula (I) as claimed in Claim 1, which compound is selected from:

6-{[(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1S)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl]carbonyl]amino}pyridine-3-carboxylic acid;

6-{[(3-[(2,2-difluoro-2-phenylethyl)oxy]-5-[(1-methylethyl)oxy]phenyl]carbonyl]amino}pyridine-3-carboxylic acid ;

20 6-{[(3-[(2-cyclopentylideneethyl)oxy]-5-[(1S)-1-methyl-2-(methyloxy)ethyl]oxy}phenyl]carbonyl]amino}pyridine-3-carboxylic acid; and

6-{[(3-[(2-cyclopentylideneethyl)oxy]-5-[(1-methylethyl)oxy]phenyl]carbonyl]amino}pyridine-3-carboxylic acid;

or a salt, solvate or pro-drug thereof.


25

10. A pharmaceutical composition comprising a compound of Formula (I) as claimed in any one of Claims 1 to 9, or a salt, solvate or prodrug thereof, together with a pharmaceutically-acceptable diluent or carrier.


30 11. A compound of Formula (I), as claimed in any one of Claims 1 to 9, or a salt, solvate or prodrug thereof, for use as a medicament.

12. A compound of Formula (I), as claimed in any one of Claims 1 to 9, or a salt, solvate or prodrug thereof, for use in the preparation of a medicament for treatment of a disease mediated through GLK, in particular type 2 diabetes.
- 5 13. A method of treating GLK mediated diseases, especially diabetes, by administering an effective amount of a compound of Formula (I), as claimed in any one of Claims 1 to 9, or a salt, solvate or prodrug thereof, to a mammal in need of such treatment.
14. The use of a compound of Formula (I), as claimed in any one of Claims 1 to 9, or salt,
10 solvate or pro-drug thereof, in the preparation of a medicament for use in the combined treatment or prevention of diabetes and obesity.
15. The use of a compound of Formula (I), as claimed in any one of Claims 1 to 9, or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the treatment or
15 prevention of obesity.
16. A method for the combined treatment of obesity and diabetes by administering an effective amount of a compound of Formula (I), as claimed in any one of Claims 1 to 9, or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.
20
17. A method for the treatment of obesity by administering an effective amount of a compound of Formula (I), as claimed in any one of Claims 1 to 9, or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.
- 25 18. A process for the preparation of a compound of Formula (I) as claimed in Claim 1, a salt, pro-drug or solvate thereof which comprises:
comprises:

(a) reaction of an acid of Formula (IIIa) or activated derivative thereof with a compound of Formula (IIIb),




Formula (IIIa)



Formula (IIIb);

5 wherein P^1 is hydrogen or a protecting group; or


(b) de-protection of a compound of Formula (IIIc),



Formula (IIIc)

wherein P^2 is a protecting group; or

10 (c) reaction of a compound of Formula (IIId) with a compound of Formula (IIIe),




Formula (IIId)

Formula (IIIe)

wherein X^1 is a leaving group and X^2 is a hydroxyl group or X^1 is a hydroxyl group and X^2 is a leaving group and wherein P^1 is hydrogen or a protecting group; or

15 (d) reaction of a compound of Formula (IIIf) with a compound of Formula (IIIg)




Formula (IIIg)

Formula (IIIf)

wherein X^3 is a leaving group and X^4 is a hydroxyl group or X^3 is a hydroxyl group and X^4 is a leaving group wherein P^1 is hydrogen or a protecting group; or

5 (e) reaction of a compound of Formula (IIIh) with a compound of Formula (IIIi),



Formula (IIIh)

Formula (IIIi);

wherein X^5 is a leaving group and wherein P^1 is hydrogen or a protecting group; and thereafter, if necessary:

- 10 i) converting a compound of Formula (I) into another compound of Formula (I);
- ii) removing any protecting groups;
- iii) forming a salt, pro-drug or solvate thereof.

INTERNATIONAL SEARCH REPORT

PCT/GB2004/005068

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D213/80 A61K31/455 A61P03/04 A61P03/10				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K A61P				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, EMBASE, MEDLINE, BIOSIS, BEILSTEIN Data, CHEM ABS Data				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	WO 03/000267 A (ASTRAZENECA AB; ASTRAZENECA UK LIMITED; HAYTER, BARRY, RAYMOND; CURRIE) 3 January 2003 (2003-01-03) cited in the application claims 1-5,7,8,13,14; examples 30-32,53,58,72,74,80,81 -----	1-18		
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.				
* Special categories of cited documents :				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top; border: none;"> *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the International filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the International filing date but later than the priority date claimed </td> <td style="width: 50%; vertical-align: top; border: none;"> *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family </td> </tr> </table>			*A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the International filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the International filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family
A document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the International filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the International filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family			
Date of the actual completion of the international search	Date of mailing of the international search report			
16 March 2005	29/03/2005			
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer vanVoorsttotVoorst, M			

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2004/005068

Box II Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 13, 16, 17
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 13, 16 and 17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of Item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/GB2004/005068

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03000267	A	03-01-2003	BR 0210711 A	20-07-2004
			CA 2451249 A1	03-01-2003
			EP 1404335 A1	07-04-2004
			WO 03000267 A1	03-01-2003
			JP 2005500312 T	06-01-2005
			MX PA03012004 A	26-03-2004
			US 2004214868 A1	28-10-2004
<hr/>				